arXiv:1701.01247v1 [cs.IT] 5 Jan 2017

Several classes of optimal ternary cyclic codes*

Lanqiang Li, Li Liu*, Shixin Zhu

School of Mathematics, Hefei University of Technology, Hefei 230009, Anhui, P.R.China

Abstract: Cyclic codes have efficient encoding and decoding algorithms over finite fields, so that they have practical applications in communication systems, consumer electronics and data storage systems. The objective of this paper is to give eight new classes of optimal ternary cyclic codes with parameters $[3^m-1, 3^m-1-2m, 4]$, according to a result on the non-existence of solutions to a certain equation over F_{3^m}. It is worth noticing that some recent conclusions on such optimal ternary cyclic codes are some special cases of our work. More importantly, three of the nine open problems proposed by Ding and Helleseth in [8] are solved completely. In addition, another one among the nine open problems is also promoted.

Keywords: Finite fields, Cyclic codes, Optimal codes, Dual codes

1. Introduction

Let F_q be the finite field with prime power $q = p^m$ elements, where p is the characteristic of the field and m is a positive integer. An $[n, k, d]$ linear code is called a cyclic code if any $(c_0, c_1, ..., c_{n-1}) \in C$ implies $(c_{n-1}, c_0, c_1, ..., c_{n-2}) \in C$. It is well known that every cyclic code of length n over F_p can be regarded as an ideal $\langle g(x)\rangle$ of the principal ring $F_p[x]/(x^n-1)$, where $g(x)|x^n-1$ is monic and has the least degree. In addition, the definition of the dual code of code C is given by

$$C^{\perp} = \{x \in F_p^n | x \cdot y = 0, \forall\ y \in C\},$$

where $x \cdot y$ is the Euclidean inner product of x and y in F_p^n.

For any codeword $c = (c_0, c_1, ..., c_{n-1}) \in C$, the Hamming weight of c is defined to be

$$w(c) = |\{i | c_i \neq 0, i = 0, 1, ..., n-1\}|.$$

Let A_i denote the number of codewords of Hamming weight i in C for $1 \leq i \leq n$, then $\{A_0, A_1, ..., A_n\}$ is said to be the weight distribution of code C. The weight enumerator of C is defined as follows:

$$A_0 + A_1 x + A_2 x^2 + ... + A_n x^n,$$

where $A_0 = 1$.

Cyclic codes over finite fields have practical applications in communication systems, consumer electronics and data storage systems, due to their efficient encoding and decoding algorithms [18, 19]. Moreover, they also have wide applications in cryptography [2, 7] and sequence design [14, 15]. All of these encourage many scholars to study cyclic codes in resent years $[1, 3-6, 21, 22, 24-30]$. In addition, it is a worthwhile subject to construct optimal cyclic codes over finite fields. For example, Carlet et al. constructed optimal ternary cyclic codes with minimum distance four, by using perfect nonlinear monomials in [2]. In 2013, Ding and Helleseth obtained some optimal ternary cyclic codes by employing almost perfect nonlinear monomials

* Corresponding author.
E-mail addresses:
lilanqiang716@126.com(Lanqiang Li), liuli-1128@163.com(Li Liu), zhushixin@hfut.edu.cn(Shixin Zhu).
This research is supported in part by the National Natural Science Foundation of China under Project 61572168, Project 11401154 and Project 11501156.

and some other monomials over F_{3^m}. They also presented nine open problems in [8]. After that, some of the open problems were solved by Li et al. in [9, 10], by discussing the non-existence of solutions to a certain equation over F_{3^m} which has been mentioned in [8]. In 2014, Zhou and Ding gave a class of optimal ternary cyclic codes with parameters $[3^m-1, 3^m-1-2m, 4]$ in [29]. Recently, Fan et al. obtained a new class of optimal ternary cyclic codes with minimum distance four. Furthermore, they also discussed the weight of the duals of the class of cyclic codes. What's more, Wang and Wu listed four classes of optimal ternary cyclic codes with parameters $[3^m-1, 3^m-1-2m, 4]$ in [12].

In this paper, we study several classes of ternary cyclic codes $C_{(1,e)}$ with generator polynomials $M(x)M_e(x)$ for suitable values of e. In Section 2, we give some basic lemmas which will be frequently used to prove our main results. Based on the non-existence of solutions to a certain equation over F_{3^m}, eight classes of ternary cyclic codes are given with parameters $[3^m-1, 3^m-1-2m, 4]$ and are thus optimal due to the Sphere Packing bound in Section 3 and Section 4. Close attention should be paid to the fact that we deal with three of the nine open problems proposed by Ding and Helleseth in [8], in Section 4. Moreover, another one among the nine open problems is promoted in Section 4.

2. Preliminaries

In this paper, we only discuss the case that $p = 3$. Let $F_{3^m}^*$ be the multiplicative cyclic group of none-zero elements of F_{3^m}. Then there must exists an element α such that $F_{3^m}^* = \langle\alpha\rangle$.

For any integer s, $0 \le s \le 3^m - 1$, the definition of 3−cyclotomic coset modulo $3^m - 1$ containing s is as follows:

$$C_s = \{s, 3s, ..., 3^{n_s-1}s\},$$

where n_s is the least positive integer such that $3^{n_s}s \equiv s(\text{mod } 3^m - 1)$. Obviously, α is a primitive $(3^m - 1)$th root of unity, then the polynomial $M_s(x) = \prod_{i\in C_s}(x - \alpha^i)$ is defined as the minimal polynomial of α^s over F_3.

Let $C_{(1,e)}$ denote the cyclic code of length $3^m - 1$ over the finite field F_3 with generator polynomial $M_1(x)M_e(x)$, where $e \notin C_1$. It is known that the cyclic codes $C_{(1,e)}$ are optimal with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$, when e are some fixed integers $[8 - 10, 12]$.

The following lemmas are very important to prove our main results, which have been given by Ding and Helleseth in [8].

Lemma 2.1. For any integer $1 \le e \le p^m - 2$ with $\gcd(e, p^m - 1) = 2$, the cardinality of the p−cyclotomic coset C_e is equal to m.

Lemma 2.2. Let $e \notin C_1$ and $|C_e| = m$. The ternary cyclic code $C_{(1,e)}$ has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ if and only if the following conditions are satisfied:
C1: e is even;
C2: the equation $(x + 1)^e + x^e + 1 = 0$ has the only solution $x = 1$ in F_{3^m}; and
C3: the equation $(x + 1)^e - x^e - 1 = 0$ has the only solution $x = 0$ in F_{3^m}.

The following three lemmas are also important and have been given by R. Lidl and H. Niederreiter in [23].

Lemma 2.3. Let q be a prime power and $g(x)$ be a nonconstant polynomial in $F_q[x]$. Then for any $f(x) \in F_q[x]$ there exist polynomials $h(x), r(x) \in F_q[x]$ such that $f(x) = g(x)h(x) + r(x)$,

where $\deg(r(x)) < \deg(g(x))$. Moreover, $\gcd(f(x), g(x)) = \gcd(g(x), r(x))$.

Lemma 2.4. For any finite field F_q and every positive integer n, where q is a prime power, the product of all monic irreducible polynomials over $F_q[x]$ whose degrees divide n is equal to $x^{q^n} - x$.

Lemma 2.5. Let q be a prime power and let $f(x)$ be an irreducible polynomial over F_q of degree n. Then $f(x) = 0$ has a root x in F_{q^n}. Furthermore, all the roots of $f(x) = 0$ are simple and are given by the n distinct elements x, x^q, x^{q^2},...,$x^{q^{n-1}}$ of F_{q^n}.

3. New optimal ternary codes with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$

In this section, we will give four new classes of optimal ternary cyclic codes $C_{(1,e)}$ with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$, by discussing the non-existence of solutions to a certain equation over F_{3^m}.

From now on, we denote the set of all nonzero squares (or nonsquares) in F_{3^m} by QR (or NQR).

3.1. The first class of optimal ternary codes

Firstly, we consider the integer e with the following form

$$e(3^s - 1) \equiv 3^t - 3^k (\text{mod } 3^m - 1), \tag{1}$$

where m, s, t, k are integers such that $0 \leq s, t \leq m - 1$ and $0 \leq k \leq t - 1$.

Lemma 3.1. Let m be a positive integer and s, t, k be integers such that $0 \leq s, t \leq m - 1$ and $0 \leq k \leq t - 1$. Then the congruence equation (1) has even solutions, if one of the following holds.
(i) m is odd and $\gcd(m, s) \mid (t - k)$.
(ii) m and $t - k$ are even, s is odd and $\gcd(m, s) \mid (t - k)$.
(iii) m, s and $m/\gcd(s, m)$ are even, $\gcd(s, m) \mid (t - k)$ and $(t - k)/\gcd(s, m)$ is even.
(iv) m and s are even, $\gcd(s, m) \mid (t - k)$ and $m/\gcd(s, m)$ are odd.

Proof. The congruence equation (1) has even solutions if and only if $(2e^{'})(3^s - 1) \equiv 3^t - 3^k (\text{mod } 3^m - 1)$ has solutions, if and only if $\gcd(2(3^s - 1), 3^m - 1) \mid (3^t - 3^k)$, i.e., $\gcd(2(3^s - 1), 3^m - 1) \mid (3^{t-k} - 1)$.

Assume that m is odd, then we have $4 \nmid (3^m - 1)$. So, $\gcd(2(3^s - 1), 3^m - 1) = \gcd(3^s - 1, 3^m - 1) = 3^{\gcd(m,s)} - 1$. Furthermore, $\gcd(2(3^s - 1), 3^m - 1) \mid (3^{t-k} - 1)$ is equivalent to $(3^{\gcd(m,s)} - 1) \mid (3^{t-k} - 1)$, i.e. $\gcd(m, s) \mid (t - k)$. Therefore, (i) is proved.

Assume that m is even, by using the similar way with the above proof, then (ii),(iii) and (iv) can be proved immediately. □

Suppose that e is an even solution of the congruence equation (1) and $\gcd(e, 3^m - 1) = 2$. According to Lemmas 2.1 and 2.2, we could know that the ternary cyclic code $C_{(1,e)}$ is optimal and has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ if and only if conditions C1, C2 and C3 are all hold. Next, we only need to prove that the equation $(x+1)^e + x^e + 1 = 0$ has the only solution $x = 1$ in F_{3^m} and the equation $(x + 1)^e - x^e - 1 = 0$ has the only solution $x = 0$ in F_{3^m}.

Therefore, we need to prove that the equations $(x+1)^e = \pm(x^e+1)$ have no solution in $F_{3^m} \setminus \{0, 1, -1\}$. By taking (3^s-1)th power on both sides of the above equations, we have

$$(x+1)^{e(3^s-1)} = (x^e+1)^{3^s-1}.$$

From (1), we have

$$(x+1)^{3^t-3^k} = (x^e+1)^{3^s-1}.$$

Furthermore, we reduce the above equation to

$$x^{3^t+e} + x^{3t} + x^e + 1 = x^{e\cdot 3^s+3^k} + x^{e\cdot 3^s} + x^{3^k} + 1.$$

By (1), we know $3^t + e = e \cdot 3^s + 3^k (\mathrm{mod}\ 3^m - 1)$. So,

$$x^{3t} + x^e = x^{e\cdot 3^s} + x^{3^k},$$

i.e.

$$x^{3^k}(x^{3^t-3^k} - 1)(x^{e-3^k} - 1) = 0.$$

It is clear that the above equation has no solution in $F_{3^m} \setminus \{0, 1, -1\}$ if $\gcd(t-k, m) = 1$ and $\gcd(e - 3^k, 3^m - 1) = 1$.

Suppose that m is even. By Lemma 3.1, we get that $t-k$ must be even if the congruence equation (1) has even solutions. This implies that $2|\gcd(t-k, m)$, i.e., $\gcd(t-k, m) \neq 1$. Therefore, there exists no optimal ternary cyclic code with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ in this subcase.

Assume that m is odd. Combining the above discussion and Lemma 3.1, we can obtain the following theorem.

Theorem 3.2. Let m be odd, $\gcd(m, s) = 1$ and e be some even solution of the congruence equation (1). Then the ternary cyclic code $C_{(1,e)}$ is optimal with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ if $\gcd(t-k, m) = 1$ and $\gcd(e - 3^k, 3^m - 1) = 1$.

proof. We know that there must be some e is an even solution of (1), due to Lemma 3.1. According to (1) and $\gcd(t-k, m) = 1$, we have $\gcd(e(3^s-1), 3^m-1) = \gcd(3^t - 3^k, 3^m - 1) = \gcd(3^{t-k} - 1, 3^m - 1) = 2$. So, $\gcd(e, 3^m - 1) = 2$, as m is odd. From Lemma 2.1, we know $|C_e| = m$. Obviously, $e \notin C_1$ as e is even. Combining Lemma 2.2 and the above discussion, the result could be derived immediately. □

Notice that $e(3^s - 1) \equiv 3^t - 1 (\mathrm{mod}\ 3^m - 1)$ is a special case of (1) with $k = 0$. In this case, we have the following corollary immediately, which was one of the main results in [12].

Corollary 3.3. Let m be odd, $\gcd(m, s) = \gcd(m, t) = 1$ and e be some solution of the congruence equation $e(3^s - 1) \equiv 3^t - 1 (\mathrm{mod}\ 3^m - 1)$. Then the ternary cyclic code $C_{(1,e)}$ is optimal with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ if $\gcd(e - 1, 3^m - 1) = 1$.

Example 1. Let $m = 3$ and $(s, t, k) = (2, 2, 1)$. Then (1) has only two solutions $e = 4$ or 17, by calculating. So, $e = 4$. Let α be the generator of $F_{3^m}^*$ with $\alpha^3 + 2\alpha + 1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[26, 20, 4]$ and is thus optimal with generator polynomial $x^6 + x^5 + 2x^4 + 2x^3 + x^2 + x + 2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 312y^{15} + 260y^{18} + 156y^{21}.$$

Example 2. Let $m = 5$ and $(s, t, k) = (4, 2, 1)$. Then (1) has only two solutions $e = 112$ or 233, i.e., $e = 112$. Let α be the generator of $F_{3^m}^*$ with $\alpha^5 + 2\alpha + 1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[242, 232, 4]$ and is thus optimal with generator polynomial $x^{10} + x^9 + 2x^6 + 2x^5 + x^4 + x + 2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 2420y^{147} + 2420y^{150} + 7260y^{153} + 4840y^{156} + 12100y^{159} + 5324y^{162}$$

$$+12100y^{165} + 4840y^{168} + 2904y^{171} + 2420y^{174} + 2420y^{177}.$$

3.2. The second class of optimal ternary codes

In this subsection, we assume that

$$e(3^s + 1) \equiv 3^t + 3^k (\text{mod } 3^m - 1), \tag{2}$$

where m, s, t, k are integers such that $0 \leq s, t, k \leq m - 1$.

Lemma 3.4. Let m be a positive integer and s, t, k be integers such that $0 \leq s, t, k \leq m - 1$. Then the congruence equation (2) has even solutions e such that $e \notin C_1$ and $|C_e| = m$, if one of the following holds.
(i) m is odd.
(ii) m is even, $\gcd(m, |t - k|) = 1$ and $m/\gcd(m, s)$ is odd.

Proof. According to the proof of Lemma 3.1, we know that first we need to prove $\gcd(2(3^s + 1), 3^m - 1) \mid (3^t + 3^k)$, i.e., $\gcd(2(3^s + 1), 3^m - 1) \mid (3^{|t-k|} + 1)$.

Assume that m is odd. We have $\gcd(2(3^s + 1), 3^m - 1) = 2$. Then, $\gcd(2(3^s + 1), 3^m - 1) \mid (3^{|t-k|} + 1)$. So the congruence equation (2) have even solutions. Furthermore, we can get $\gcd(e(3^s + 1), 3^m - 1) = \gcd(3^t + 3^k, 3^m - 1) = 2$. Then $\gcd(e, 3^m - 1) = 2$ as e is even. From Lemma 2.1, we have $|C_e| = m$. It is clear that $e \notin C_1$ as e is even. Therefore, this completes the proof of (i).

Suppose that m is even. As $m/\gcd(m, s)$ are odd, we can get $\gcd(3^s + 1, 3^m - 1) = 2$. Furthermore, we have $\gcd(2(3^s + 1), 3^m - 1) = 4$, as m is even. According to $\gcd(m, |t - k|) = 1$, we have $\gcd(e(3^s + 1), 3^m - 1) = \gcd(3^t + 3^k, 3^m - 1) = 4$, which shows that (2) has even solutions. If $\gcd(e, 3^m - 1) = 4$, then $8 \mid \gcd(e(3^s + 1), 3^m - 1)$ as m is even, which is a contradiction. Then, it is easy to get that $\gcd(e, 3^m - 1) = 2$. So, we obtain $|C_e| = m$, by Lemma 2.1. Clearly, $e \notin C_1$ as e is even. □

From Lemma 3.4, we can get the following theorem.

Theorem 3.5. Let m be a positive integer, s, t, k be integers such that $0 \leq s, t, k \leq m - 1$ and e be some even solution of (2). Then the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ and is optimal if one of the following holds.
(i) m is odd, $\gcd(3^t - e, 3^m - 1) = 1$ and $\gcd(e - 3^k, 3^m - 1) = 1$.
(ii) m is even, $\gcd(m, |t - k|) = 1$, $m/\gcd(m, s)$ is odd, $\gcd(3^t - e, 3^m - 1) = 1$ and $\gcd(e - 3^k, 3^m - 1) = 1$.

Proof. Similar to the discussion of Subsection 3.1, we could obtain that conditions C2 and C3 in Lemma 2.2 are satisfied if $\gcd(3^t - e, 3^m - 1) = 1$ and $\gcd(e - 3^k, 3^m - 1) = 1$. Combining

Lemmas 2.2 and 3.4, we could get Theorem 3.5 immediately. □

Note that $e(3^s+1) \equiv 3^t+1 (\text{mod } 3^m-1)$ is a special case of (1) with $k=0$. In this case, it has been shown in [12, Theorem 2], which is covered by Theorem 3.5. Moreover, it is worth noticing that more optimal ternary cyclic codes can be found than [12].

Example 3. Let $m=7$ and $(s,t,k)=(4,2,1)$. Then (2) has only two solutions $e=480$ or 1573. So, $e=480$. Let α be the generator of $F_{3^m}^*$ with $\alpha^7+2\alpha^2+1=0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[2186, 2172, 4]$ and is optimal with generator polynomial $x^{14}+x^{13}+2x^{12}+x^{11}+2x^{10}+x^7+x^3+2x^2+x+2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1+153020y^{1404}+1040536y^{1431}+2513900y^{1458}+922492y^{1485}+153020y^{1512}.$$

Example 4. Let $m=4$ and $(s,t,k)=(0,2,1)$. Then (2) has only two even solutions $e=6$ or 46. Let α be the generator of $F_{3^m}^*$ with $\alpha^4-\alpha^3+2=0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[80, 72, 4]$ and is optimal with generator polynomial $x^8+x^7+2x^6+2x^5+2x^2+2(e=6)$ or $x^8+2x^5+x^3+2x^2+2(e=46)$. The dual of $C_{(1,e)}$ for the cases of $e=6$ or $e=46$ has weight enumerator

$$1+1320y^{48}+2400y^{51}+80y^{54}+1920y^{57}+840y^{60}.$$

Example 5. Let $m=6$ and $(s,t,k)=(0,2,1)$. Then (2) has only two even solutions $e=6$ or 370. Let α be the generator of $F_{3^m}^*$ with $\alpha^4-\alpha^3+2=0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[728, 716, 4]$ and is optimal with generator polynomial $x^{12}+x^{11}+2x^{10}+x^9+x^8+x^7+2x^6+x^5+x^3+x^2+2x+2(e=6)$ or $x^{12}+2x^{11}+2x^{10}+2x^9+x^8+x^6+2x^5+x^4+2x^3+x^2+2x+2(e=370)$. The dual of $C_{(1,e)}$ for the cases of $e=6$ or $e=370$ has weight enumerator

$$1+95004y^{468}+183456y^{477}+728y^{486}+170352y^{495}+81900y^{504}.$$

3.3. The third class of optimal ternary codes

In this subsection, we consider the following form on e,

$$e \equiv (3^{m-1}-1)/2+3^h (\text{mod } 3^m-1), \tag{3}$$

where h is an integer such that $0 \le h \le m-1$.

Lemma 3.6. Let e be given by (3). Then e is even and $|C_e| = m$ if m is even and $\gcd(m, h+1)=1$.

Proof. Obviously, e is even if m is even. Notice that $\gcd(3e, 3^m-1) = \gcd(\frac{3^m-1}{2}+3^{h+1}-1, 3^m-1) = \gcd(3^{h+1}-1, 3^m-1) = 2$, as $\gcd(m, h+1)=1$. So, $\gcd(e, 3^m-1)=2$. From Lemma 2.1, we have $|C_e|=m$. □

Theorem 3.7. Let m be even and e has the form as (3). Then the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m-1, 3^m-1-2m, 4]$ and is optimal if $\gcd(m, h+1)=1$ and $\gcd(3^{h+1}-2, 3^m-1)=1$.

Proof. From Lemma 3.6, we know that it is enough to prove that $(x+1)^e \pm (x^e+1) = 0$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$. By taking 3th power on both sides of the above equation, we could get $(x+1)^{3e} \pm (x^{3e}+1) = 0$. Obviously, we know that $(x+1)^{3e} \pm (x^{3e}+1) = 0$ and $(x+1)^e \pm (x^e+1) = 0$ have the same solutions. Next, we could discuss the solutions of the equation $(x+1)^{3e} \pm (x^{3e}+1) = 0$ as follows:

(1)If $x, x+1 \in$ QR or $x \in$ QR and $x+1 \in$ NQR. Since $3e = \frac{3^m-1}{2} + 3^{h+1} - 1$, $(x+1)^{3e} \pm (x^{3e}+1) = 0$ is equivalent to $(x+1)^{3^{h+1}-1} \pm (x^{3^{h+1}-1}+1) = 0$ or $-(x+1)^{3^{h+1}-1} \pm (x^{3^{h+1}-1}+1) = 0$. Multiplying both sides of the above two equations by $x(x+1)$, we could obtain that $x(x^{3^{h+1}-1}-1)(x-1) = 0$ or $x^2(x^{3^{h+1}-2}+1) = 0$ have no solution in $F_{3^m} \setminus \{0, 1, -1\}$ if $\gcd(m, h+1) = 1$ and $\gcd(3^{h+1}-2, 3^m-1) = 1$.

(2)If $x, x+1 \in$ NQR or $x \in$ NQR and $x+1 \in$ QR. In this case, using the similar method in (1), we can obtain that $(x+1)^{3e} \pm (x^{3e}+1) = 0$ is equivalent to $(x+1)^2((x+1)^{3^{h+1}-2}-1) = 0$ or $x^{3^{h+1}+1} - x^{3^{h+1}} + x^2 = 0$. It is easy to see that the equation $(x+1)^2((x+1)^{3^{h+1}-2}-1) = 0$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$ if $\gcd(3^{h+1}-2, 3^m-1) = 1$.

Moreover, suppose that $x_0 \in F_{3^m} \setminus \{0, 1, -1\}$ is some solution of the equation $x^{3^{h+1}+1} - x^{3^{h+1}} + x^2 = 0$. Then we know that x_0 must also be a solution of the differential of the equation, i.e., $x_0^{3^{h+1}} - x_0 = 0$. This is a contradiction with $\gcd(m, h+1) = 1$ and $x_0 \in F_{3^m} \setminus \{0, 1, -1\}$. □

Note that $e = (3^{m-1}-1)/2 + 1$ if $h = 0$ in (3). Furthermore, by Theorem 3.7, we could obtain the following corollary.

Corollary 3.8. Let $e = (3^{m-1}-1)/2 + 1$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ and is optimal if m is even.

Proof. Obviously, $e = (3^{m-1}-1)/2 + 1$ is a special case of (3) with $h = 0$. Furthermore, we have $\gcd(m, h+1) = 1$ and $\gcd(3^{h+1}-2, 3^m-1) = 1$, when $h = 0$. According to Theorem 3.7, we can get the result immediately. □

Example 6. Let $(m, h) = (4, 0)$, i.e., the case in Corollary 3.8. Then $e = 14$. Let α be the generator of $F_{3^m}^*$ with $\alpha^4 + 2\alpha^3 + 2 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[80, 72, 4]$ and is optimal with generator polynomial $x^8 + 2x^5 + x^3 + 2x^2 + 2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 1320y^{48} + 2400y^{51} + 80y^{54} + 1920y^{57} + 840y^{60}.$$

Example 7. Let $(m, h) = (6, 0)$, then $e = 122$. Let α be the generator of $F_{3^m}^*$ with $\alpha^6 + 2\alpha^4 + \alpha^2 + 2\alpha + 2 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[728, 716, 4]$ and is optimal with generator polynomial $x^{12} + 2x^{11} + 2x^{10} + 2x^9 + x^8 + x^6 + 2x^5 + x^4 + 2x^3 + x^2 + 2x + 2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 95004y^{468} + 183456y^{477} + 728y^{486} + 170352y^{495} + 81900y^{504}.$$

Example 8. Let $(m, h) = (6, 4)$, then $e = 202$. Let α be the generator of $F_{3^m}^*$ with $\alpha^6 + 2\alpha^4 + \alpha^2 + 2\alpha + 2 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[728, 716, 4]$ and is optimal with generator polynomial $x^{12} + 2x^{10} + x^9 + x^8 + 2x^5 + x^4 + x^3 + 2x^2 + 2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 1456y^{450} + 8736y^{456} + 8736y^{462} + 8736y^{465} + 22204y^{468} + 39312y^{471} + 34944y^{474}$$

$$+52416y^{477} + 26208y^{480} + 65520y^{483} + 50232y^{486} + 17472y^{489} + 26208y^{492}$$
$$+52416y^{495} + 56784y^{498} + 17472y^{501} + 12012y^{504} + 21840y^{507} + 8736y^{519}.$$

3.4. The fourth class of optimal ternary codes

In this subsection, we consider the following form on e,

$$e \equiv (3^{m-1} - 1)/2 + 3^h - 1 (\text{mod } 3^m - 1), \tag{4}$$

where h is an integer such that $0 \leq h \leq m-1$.

Obviously, we know that e is even only if m is odd.

Lemma 3.9. Let e be an integer and be given by (4). Then e is even and $|C_e| = m$ if m is odd prime or m is odd and $\gcd(3^{h+1} - 4, 3^m - 1) = 1$.

Proof. It is clear that e is even if m is odd. If m is odd prime, then $|C_e| = m$ as $e \neq 0$. Moreover, note that $\gcd(3e, 3^m - 1) = \gcd(\frac{3^m-1}{2} + 3^{h+1} - 4, 3^m - 1) = 2 \cdot \gcd(3^{h+1} - 4, 3^m - 1) = 2$, if $\gcd(3^{h+1} - 4, 3^m - 1) = 1$. So, we have $\gcd(e, 3^m - 1) = 2$. By Lemma 2.1, we have $|C_e| = m$. □

Theorem 3.10. Let m be odd and e be given by (4). Then the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ and is optimal if m is odd prime and $\gcd(3^{h+1} - 5, 3^m - 1) = 2$ or $\gcd(3^{h+1} - 4, 3^m - 1) = 1$ and $\gcd(3^{h+1} - 5, 3^m - 1) = 2$.

Proof. Obviously, e is even and $|C_e| = m$ from Lemma 3.9. Similar to the proof of Theorem 3.7, we only need to discuss the solutions of the equation $(x+1)^{3e} \pm (x^{3e} + 1) = 0$ as follows:

(i) $x, x+1 \in$ QR or $x \in$ QR and $x+1 \in$ NQR. As $3e = \frac{3^m-1}{2} + 3^{h+1} - 4$, the equation $(x+1)^{3e} \pm (x^{3e} + 1) = 0$ can be simplified to $x^{3^{h+1}+1} - x^{3^{h+1}-1} - x^{3^{h+1}-3} - x^{3^{h+1}-4} - x^4 - x^3 - x + 1 = 0$ or $x^{3^{h+1}-1} + x^{3^{h+1}-3} + x^{3^{h+1}-4} + x^4 + x^3 + x = 0$.

Let $f(x) = x^{3^{h+1}+1} - x^{3^{h+1}-1} - x^{3^{h+1}-3} - x^{3^{h+1}-4} - x^4 - x^3 - x + 1 \in F_{3^m}[x]$. Moreover, we suppose that $x_0 \in F_{3^m} \setminus \{0, 1, -1\}$ is some root of $f(x)$. Obviously, x_0 is a 3-heavy root of $f(x)$. Therefore, x_0 is also a root of $f''(x) = x^{3^{h+1}-6}(x+1)^3$. However, notice that $f''(x) = x^{3^{h+1}-6}(x+1)^3$ only has two solutions $x = 0, -1$, which shows that $f(x)$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$.

Let $f(x) = x^{3^{h+1}-1} + x^{3^{h+1}-3} + x^{3^{h+1}-4} + x^4 + x^3 + x$, then $f'(x) = (x^{3^{h+1}-5} - 1)(x+1)^3$. Furthermore, we know that $f'(x)$ only has two solutions $x = 1, -1$ in F_{3^m}, as $\gcd(3^{h+1} - 5, 3^m - 1) = 2$. Since every root of $f(x)$ is 3-heavy root, we could deduce that $f(x)$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$ when $\gcd(3^{h+1} - 5, 3^m - 1) = 2$.

(ii) $x, x+1 \in$ NQR or $x \in$ NQR and $x+1 \in$ QR. In this case, $(x+1)^{3e} \pm (x^{3e} + 1) = 0$ can be reduced to $x^{3^{h+1}} - x^{3^{h+1}-1} - x^{3^{h+1}-3} - x^{3^{h+1}-4} + x^4 + x^3 + x = 0$ or $x^{3^{h+1}-1} + x^{3^{h+1}-3} + x^{3^{h+1}-4} - x^4 - x^3 - x + 1 = 0$.

Let $f(x) = x^{3^{h+1}} - x^{3^{h+1}-1} - x^{3^{h+1}-3} - x^{3^{h+1}-4} + x^4 + x^3 + x$, then $f'(x) = (x^{3^{h+1}-5} + 1)(x+1)^3$. Due to m is odd, we have $\gcd(2(3^{h+1} - 5), 3^m - 1) = 2$. This implies that $f'(x)$ only has two solutions $x = 1, -1$ in F_{3^m}. So, we can deduce that $f(x)$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$.

Let $f(x) = x^{3^{h+1}-1} + x^{3^{h+1}-3} + x^{3^{h+1}-4} - x^4 - x^3 - x + 1$, then $f''(x) = x^{3^{h+1}-6}(x+1)^3$. Similar to (i), we can get that $f(x)$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$. □

Note that $e = (3^{m-1}-1)/2$ if $h = 0$ in (5). In this special case, we can obtain the following corollary.

Corollary 3.11. Let $e = (3^{m-1}-1)/2$. Then the ternary cyclic code $C_{(1,e)}$ is optimal with parameters $[3^m-1, 3^m-1-2m, 4]$ if m is odd.

Proof. If $h = 0$, then $\gcd(3^{h+1}-4, 3^m-1) = 1$ and $\gcd(3^{h+1}-5, 3^m-1) = 2$. So, we can get the result immediately, according to Theorem 3.10. □

Example 9. Let $(m, h) = (5, 0)$, then $e = 40$. Let α be the generator of $F_{3^m}^*$ with $\alpha^5+2\alpha+1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[242, 232, 4]$ and is optimal with generator polynomial $x^{10}+x^9+2x^6+2x^5+x^4+x+2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1+2420y^{147}+2420y^{150}+7260y^{153}+4840y^{156}+12100y^{159}+5324y^{162}$$
$$+12100y^{165}+4840y^{168}+2904y^{171}+2420y^{174}+2420y^{177}.$$

Example 10. Let $(m, h) = (5, 3)$ then $e = 66$. Let α be the generator of $F_{3^m}^*$ with $\alpha^5+2\alpha+1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[242, 232, 4]$ and is optimal with generator polynomial $x^{10}+x^9+2x^8+x^7+2x^6+2x^5+2x^4+x^3+x^2+x+2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1+132y^{132}+7260y^{150}+7260y^{153}+9680y^{159}+14762y^{162}+14520y^{168}+4840y^{171}+484y^{186}+110y^{198}.$$

Example 11. Let $(m, h) = (3, 2)$, then $e = 12$. Let α be the generator of $F_{3^m}^*$ with $\alpha^3+2\alpha+1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[26, 20, 4]$ and is optimal with generator polynomial $x^6+x^5+2x^4+2x^3+x^2+x+2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1+312y^{15}+260y^{18}+156y^{21}.$$

4. Four open problems about the ternary cyclic codes $C_{(1,e)}$

In this section, we will explicitly deal with three open problems among nine open problems about the ternary cyclic codes $C_{(1,e)}$ which have been proposed by Ding and Helleseth in [8]. Moreover, another one among nine open problems is promoted. So, four new classes of optimal ternary cyclic codes $C_{(1,e)}$ with parameters $[3^m-1, 3^m-1-2m, 4]$ are obtained.

4.1. Solving the first open problem

In this subsection, we discuss the following form on e which has been given as open problem 7.15 by Ding and Helleseth in [8],

$$e \equiv (3^{m-1}-1)/2 + 3^h + 1 (\mathrm{mod}\ 3^m-1), \tag{5}$$

where h is an integer such that $0 \le h \le m-1$.

Open problem 1.(see [8, open problem 7.15]) Let $e = (3^{m-1}-1)/2+3^h+1$, where $0 \le h \le m-1$. What are the conditions on m and h under which the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m-1, 3^m-1-2m, 4]$?

In order to solve this problem, the following lemma is necessary.

Lemma 4.1. Let e be an integer with the form (5). Then e is even and $|C_e| = m$ if m is odd prime or m is odd and $\gcd(3^{h+1}+2, 3^m-1) = 1$.

Proof. Obviously, e is even if m is odd. If m is odd prime, then $|C_e| = m$ as $e \neq 0$. Moreover, notice that $\gcd(3e, 3^m-1) = \gcd(\frac{3^m-1}{2} + 3^{h+1} + 2, 3^m-1) = 2$, if $\gcd(3^{h+1}+2, 3^m-1) = 1$. So $\gcd(e, 3^m-1) = 2$. From Lemma 2.1, we have $|C_e| = m$. □

Theorem 4.2. Let m be odd and e has the form as (5). Then the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m-1, 3^m-1-2m, 4]$ and is optimal if m is odd prime with $\gcd(m, h+1) = 1$ or $\gcd(m, h+1) = 1$ and $\gcd(3^{h+1}+2, 3^m-1) = 1$.

Proof. Clearly, e is even and $|C_e| = m$ by Lemma 4.1. Using same discussion in Theorem 3.7, it is clear that we could discuss the solutions of the equation $(x+1)^{3e} \pm (x^{3e}+1) = 0$ as follows:

(i) $x, x+1 \in$ QR or $x \in$ QR and $x+1 \in$ NQR. Since $3e = \frac{3^m-1}{2} + 3^{h+1} + 2$, $(x+1)^{3e} \pm (x^{3e}+1) = 0$ is equivalent to $x^{3^{h+1}+2} + x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x + 1 = 0$ or $x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x = 0$.

Let $f(x) = x^{3^{h+1}+2} + x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x + 1$, then the differential $f' = -(x-1)^{3^{h+1}+1} - 1$ of the polynomial $f(x)$. Let $f'(x) = 0$, then $(x-1)^{2(3^{h+1}+1)} = 1$. As m is odd, we have $\gcd(2(3^{h+1}+1), 3^m-1) = 2$. So, the equation $(x-1)^{2(3^{h+1}+1)} = 1$ only has two solutions $x = 0, -1$ which implies that $f(x)$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$.

If $x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x = 0$, i.e., $x(x^{3^{h+1}-1} - 1)(x-1) = 0$, then the equation has no solution in $F_{3^m} \setminus \{0, 1, -1\}$ when $\gcd(3^m-1, 3^{h+1}-1) = 2$, i.e., $\gcd(m, h+1) = 1$.

(ii) $x, x+1 \in$ NQR or $x \in$ NQR and $x+1 \in$ QR. In this case, we can obtain that $(x+1)^{3e} \pm (x^{3e}+1) = 0$ is equivalent to $x^{3^{h+1}+2} + x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x = 0$ or $x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x + 1 = 0$.

Let $f(x) = x^{3^{h+1}+2} + x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x$, then $f' = -(x-1)^{3^{h+1}+1} - 1$. Clearly, $f(x)$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$, by (i).

Let $f(x) = x^{3^{h+1}+1} - x^{3^{h+1}} - x^2 + x + 1$, then $f'(x) = x^{3^{h+1}} + x + 1$. Furthermore, we have $f'' = 1$ in $F_{3^m}(x)$ which means $f(x)$ has no 3−heavy solution in $F_{3^m} \setminus \{0, 1, -1\}$, i.e., $f(x)$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$. □

Clearly, $e = (3^{m-1}-1)/2 + 2$ is a special case of (5) with $h = 0$. So, we can obtain the following corollary immediately.

Corollary 4.3. Let $e = (3^{m-1}-1)/2 + 2$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m-1, 3^m-1-2m, 4]$ and is optimal if m is odd.

Remark 1. For $m \in \{3, 5, 7, 9, 11\}$, the results of Theorem 4.2 are positive and verified by Magma. Some concrete examples are listed as follows.

Example 12. Let $(m, h) = (3, 0)$, then $e = 6$. Let α be the generator of $F_{3^m}^*$ with $\alpha^3 + 2\alpha + 1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[26, 20, 4]$ and is optimal with generator polynomial $x^6 + x^5 + 2x^3 + 2x + 2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 312y^{15} + 260y^{18} + 156y^{21}.$$

Example 13. Let $(m,h)=(5,2)$, then $e=50$. Let α be the generator of $F_{3^m}^*$ with $\alpha^5+2\alpha+1=0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[242,232,4]$ and is optimal with generator polynomial $x^{10}+2x^8+2x^7+2x^6+x^4+2x^2+x+2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1+2420y^{144}+12100y^{153}+34364y^{162}+7744y^{171}+2420y^{180}.$$

Example 14. Let $(m,h)=(5,3)$, then $e=68$. Let α be the generator of $F_{3^m}^*$ with $\alpha^5+2\alpha+1=0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[242,232,4]$ and is optimal with generator polynomial $x^{10}+2x^9+2x^8+2x^6+x^5+2x^3+a+2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1+2420y^{144}+12100y^{153}+34364y^{162}+7744y^{171}+2420y^{180}.$$

4.2. Solving the second open problem

In this subsection, we settle a problem which has been proposed as open problem 7.13 by Ding and Helleseth in [8] as follows:

Open problem 2.(see [8, open problem 7.13]) Let $e=3^h+5$, where $0\leq h\leq m-1$. Let $m\geq 5$ and let m be a prime. Is it true that the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m-1,3^m-1-2m,4]$ for all h with $0\leq h\leq m-1$?

In order to solve this problem, we must first to prove that the following lemma hold.

Lemma 4.4. Let $e=3^h+5$, where $0\leq h\leq m-1$. Let $m\geq 5$ and let m be a prime. Then $e\notin C_1$ is even and $|C_e|=m$.

Proof. Clearly, $e=3^h+5$ is even for any h with $0\leq h\leq m-1$. So $e\notin C_1$. It is easy to get $|C_e|=m$, as m is a prime and $e\neq 0$. □

Theorem 4.5. Let $e=3^h+5$, where $0\leq h\leq m-1$. Let $m\geq 5$ and let m be a prime. Then the ternary cyclic code $C_{(1,e)}$ is optimal and has parameters $[3^m-1,3^m-1-2m,4]$ for all h with $0\leq h\leq m-1$.

Proof. Clearly, it is enough to prove that conclusions C2 and C3 in Lemma 2.2 are satisfied, from Lemma 4.4.

Firstly, we need to prove $(x+1)^e+x^e+1=0$ has no solution in $F_{3^m}\setminus\{0,1,-1\}$. By taking 3^{m-h}th power on both sides of the above equation, we obtain $(x+1)^{5\cdot 3^{m-h}+1}+x^{5\cdot 3^{m-h}+1}+1=0$. So, $(x+1)^{5\cdot 3^{m-h}+1}+x^{5\cdot 3^{m-h}+1}+1=0$ and $(x+1)^e+x^e+1=0$ have the same solutions. Let $f(x)=(x+1)^{5\cdot 3^{m-h}+1}+x^{5\cdot 3^{m-h}+1}+1$, then we know that every root of $f(x)$ must be 3^{m-h}-heavy root, according to the above discussion.

Assume that there exists some $x_0\in F_{3^m}\setminus\{0,1,-1\}$ is a root of $f(x)$. Then x_0 is also a root of $f^{'}(x)=(-x^5-x^4+x^3+x^2-x+1)^{3^{m-h}}$. However, it is easy to see that $(-x^5-x^4+x^3+x^2-x+1)=(1-x)(x^4-x^3+x^2+1)$. Next, denote $g(x)=x^4-x^3+x^2+1$. According to Lemma 2.3, we have that $\gcd(g(x),x^{3^2}-x)=1$, $\gcd(g(x),x^{3^3}-x)=1$ and $\gcd(g(x),x^{3^4}-x)=g(x)$. Then $f(x)$ has the only root $x=1$ in F_{3^m} if and only if $m\not\equiv 0(\mathrm{mod}\ 4)$, by Lemma 2.5. Since m is a prime, then $(x+1)^{5\cdot 3^{m-h}+1}+x^{5\cdot 3^{m-h}+1}+1=0$ has the only solution $x=1$ which is a contradiction with x_0 is a root of $f(x)$. Thus, $(x+1)^e+x^e+1=0$ has no solution in $F_{3^m}\setminus\{0,1,-1\}$.

Secondly, similar to the above discussion, we could prove that $(x+1)^e - x^e - 1 = 0$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$. □

Remark 2. For $m \in \{3, 5, 7, 11, 13, 17\}$, the ternary cyclic codes $C_{(1,e)}$ are all optimal with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$, where $e = 3^h + 5$ with $0 \leq h \leq m - 1$, which have been confirmed by Magma in [8].

4.3. Solving the third open problem

In this subsection, we consider the open problem 7.14 in [8] which have been given by Ding and Helleseth as follows:

Open problem 3.(see [8, open problem 7.14]) Let $e = 3^h + 13$, where $0 \leq h \leq m - 1$. Let m be an odd prime. What are the conditions on h under which the ternary cyclic codes $C_(1, e)$ has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$?

The following Lemma is necessary for solving the open problem 3.

Lemma 4.6. Let $e = 3^h + 13$, where $0 \leq h \leq m - 1$. Let m be an odd prime. Then $e \notin C_1$ is even and $|C_e| = m$.

Proof. Similar to the proof of Lemma 4.4, we could obtain the lemma immediately. □

Theorem 4.7. Let m be an odd prime with $m \not\equiv 0 (\text{mod } 3)$ and $e = 3^h + 13$ with $0 \leq h \leq m - 1$. Then the ternary cyclic code $C_{(1,e)}$ is optimal and has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ for all h with $0 \leq h \leq m - 1$.

Proof. According to Lemma 4.6, we only need to prove that conclusions C2 and C3 hold.

In the first place, we verify that $(x+1)^e + (x^e + 1) = 0$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$. Taking 3^{m-h}th power on both sides of the above equation, we can get $(x+1)^{13 \cdot 3^{m-h}+1} + x^{13 \cdot 3^{m-h}+1} + 1 = 0$. Let $f(x) = (x+1)^{13 \cdot 3^{m-h}+1} + x^{13 \cdot 3^{m-h}+1} + 1$, then any root of $f(x)$ is 3^{m-h}-heavy root which means that all the roots of $f(x)$ are also the roots of $f^{'}(x) = (x^{13} - x^{12} - x^{10} - x^9 - x^4 - x^3 - x - 1)^{3^{m-h}}$. we denote $g(x) = x^{13} - x^{12} - x^{10} - x^9 - x^4 - x^3 - x - 1$. According to Lemma 2.3, we could get that $\gcd(g(x), x^3 - x) = x - 1$, $\gcd(g(x), x^{3^2} - x) = x - 1$, $\gcd(g(x), x^{3^3} - x) = g(x)$ and $\gcd(g(x), x^{3^4} - x) = x - 1$. Then, Combining Lemma 2.4 and $\deg(g(x)) = 13$, we know that $g(x)$ has the factor $(x - 1)$ and four cubic irreducible factors. Furthermore, we have $g(x) = (x-1)(x^3-x^2-x-1)(x^3+x^2+x-1)(-x^3-x^2+x-1)(-x^3+x^2-1)$. From Lemma 2.5, it is easy to get that $g(x)$ (i.e., $f^{'}(x)$) has the only root $x = 1$ if and only if $m \not\equiv 0 (\text{mod } 3)$ which shows that $f(x)$ has no root in $F_{3^m} \setminus \{0, 1, -1\}$. So, the equation $(x+1)^e + (x^e + 1) = 0$ has no solution in $F_{3^m} \setminus \{0, 1, -1\}$.

In the second place, using the similar method with the above proof, we could get $(x + 1)^{13 \cdot 3^{m-h}+1} - x^{13 \cdot 3^{m-h}+1} - 1 = 0$. Let $f(x) = (x+1)^{13 \cdot 3^{m-h}+1} - x^{13 \cdot 3^{m-h}+1} - 1$. Then, all the roots of $f(x)$ are also the roots of $f^{'}(x) = (x^{12} + x^{10} + x^9 + x^4 + x^3 + x + 1)^{3^{m-h}}$. Denote $g(x) = x^{12} + x^{10} + x^9 + x^4 + x^3 + x + 1$. According to Lemma 4.2, we have that $\gcd(g(x), x^3 - x) = 1$, $\gcd(g(x), x^{3^2} - x) = 1$ and $\gcd(g(x), x^{3^3} - x) = g(x)$. Combining Lemma 2.4 and $\deg(g(x)) = 12$, we know that $g(x)$ has four cubic irreducible factors. Then, $f(x)$ has

no root in $F_{3^m} \setminus \{0, 1, -1\}$ if and only if $m \not\equiv 0 (\mathrm{mod}\ 3)$. This completes the proof. □

Remark 3. For $m \in \{3, 5, 7, 11\}$, the result of Theorem 4.7 is positive and confirmed by Magma. Some specific examples are given as follows.

Example 15. Let $(m, h) = (5, 0)$ then $e = 14$. Let α be the generator of $F_{3^m}^*$ with $\alpha^5+2\alpha+1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[242, 232, 4]$ and is optimal with generator polynomial $x^{10}+x^9+x^8+2x^7+2x^5+2x^3+x^2+2x+2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 21780y^{153} + 19844y^{162} + 17424y^{171}.$$

Example 16. Let $(m, h) = (5, 2)$, then $e = 22$. Let α be the generator of $F_{3^m}^*$ with $\alpha^5+2\alpha+1 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[242, 232, 4]$ and is optimal with generator polynomial $x^{10} + x^9 + 2x^8 + x^7 + 2x^6 + 2x^5 + 2x^4 + x^3 + x^2 + x + 2$. The dual of $C_{(1,e)}$ has weight enumerator

$$1+132y^{132}+7260y^{150}+7260y^{153}+9680y^{159}+14762y^{162}+14520y^{168}+4840y^{171}+484y^{186}+110y^{198}.$$

4.4. Promoting the fourth open problem

In this subsection, we discuss the open problem 7.12 in [8] which have been given by Ding and Helleseth as follows:

Open problem 4.(see [8, open problem 7.12]) Let $e = 3^h + 5$, where $2 \le h \le m - 1$. Let $m \ge 5$ and let m be even. Is it true that the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ if one of the following conditions is met?

1) $m \equiv 0 (\mathrm{mod}\ 4), m \ge 4$ and $h = m/2$.
2) $m \equiv 2 (\mathrm{mod}\ 4), m \ge 6$ and $h = (m + 2)/2$.

Theorem 4.8. Let $e = 3^h + 5$, where $2 \le h \le m - 1$. Let $m \ge 5$ and let m be even. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ if $m \equiv 2 (\mathrm{mod}\ 4), m \ge 6$ and $h = (m + 2)/2$.

Proof. Obviously, $e \notin C_1$ as e is even. Let $m = 4k + 2$ for some positive integer k, then $h = 2k + 2$. Note that $3^m - 1 = 3^{4k+2} - 1 = (3^{2k+1} - 1)(3^{2k+1} + 1)$. Furthermore, we have $\gcd(3^h + 5, 3^{2k+1} - 1) = \gcd(3(3^{2k+1} - 1) + 8, 3^{2k+1} - 1) = 2$ and $\gcd(3^h + 5, 3^{2k+1} + 1) = \gcd(3(3^{2k+1}+1)+2, 3^{2k+1}+1) = 2$. Since $4 \not| (3^h+5)$, we could obtain that $\gcd(3^h+5, 3^m-1) = 2$. By Lemma 2.1, we know that $|C_e| = m$. According to the proof of Theorem 4.5, we know that conditions C2 and C3 are satisfied if $m \not\equiv 0 (\mathrm{mod}\ 4)$. So, this completes the proof. □

Remark 4. We spare no efforts to solve the open problem 4 completely. However, we could only prove that the ternary cyclic codes $C_{(1,e)}$ are optimal with parameters $[3^m - 1, 3^m - 1 - 2m, 4]$ if condition 2) hold. As for condition 1), we could not prove the ternary cyclic codes $C_{(1,e)}$ are optimal, which means that new techniques are required.

Example 17. Let $m = 6$. Then $h = 4$ and $e = 86$. Let α be the generator of $F_{3^m}^*$ with $\alpha^6 + 2\alpha^4 + \alpha^2 + 2\alpha + 2 = 0$. Then the ternary cyclic code $C_{(1,e)}$ has parameters $[728, 716, 4]$ and is optimal with generator polynomial $x^{12} + x^{11} + 2x^{10} + x^9 + x^7 + x^5 + 2x^4 + 2x^3 + 2x^2 + 2x + 2$.

The dual of $C_{(1,e)}$ has weight enumerator

$$1 + 8736y^{450} + 8736y^{459} + 49868y^{468} + 133952y^{477} + 171080y^{486}$$
$$+91728y^{495} + 39676y^{504} + 26208y^{513} + 1456y^{522}.$$

5. Conclusions

In this paper, eight new classes of optimal ternary cyclic codes with parameters $[3^m-1, 3^m-1-2m, 4]$ were studied, by discussing the non-existence of solutions to a certain equation over F_{3^m}. The first two classes of optimal ternary cyclic codes are more general cases of some results in [12]. Furthermore, it is worth noticing that some recent conclusions on such optimal ternary cyclic codes are some special cases of our work. Moreover, the other two new classes of optimal ternary cyclic codes are also important in Section 3. More importantly, we solve three of the nine open problems proposed by Ding and Helleseth in [8] completely. In addition, another one among the nine open problems is also promoted in Section 4. It is easy to see that most of the duals of the optimal ternary codes listed in this paper have few nonzero weights. So, it would be an interesting problem to determine the weight distribution of these dual codes.

References

[1] H.Q. Dinh, On the linear odering of some classess of negacyclic and cyclic codes and their distance distributions, Finite Fields Appl. 14 (2008) 22-40.

[2] C. Carlet, C. Ding, J. Yuan, Linear codes from highly nonlinear functions and their secret sharing schemes, IEEE Trans. Inf. Theory 51 (6) (2005) 2089-2102.

[3] C. Ding, S. Ling, A q-polynomial approach to cyclic codes, Finite Fields Appl. 20 (3) (2013) 1-14.

[4] Y. Jia, S. Ling, C. Xing, On self-dual cyclic codes over finite fields, IEEE Trans. Inf. Theory 57 (2011) 2243-2251.

[5] C. Ding, Cyclic codes from some monomials and trinomials, SIAM J. Discrete Math. 27 (4) (2013) 1977-1994.

[6] X. Kai, S. Zhu, On cyclic self-dual codes, Appl. Algebra Engrg. Comm. Comput. 19 (2008) 509-525.

[7] C. Ding, X. Wang, A coding theory construction of new systematic authentication codes, Theor. Comput. Sci. 330 (1) (2005) 81-99.

[8] C. Ding, T. Helleseth, Optimal ternary cyclic codes from monomials, IEEE Trans. Inf. Theory 59 (9) (2013) 5898-5904.

[9] N. Li, C. Li, T. Helleseth, C. Ding, X.H. Tang, Optimal ternary cyclic codes with minimum distance four and five, Finite Fields Appl. 30 (2014) 100-120.

[10] N. Li, Z. Zhou, T. Helleseth, On a conjecture about a class of optimal ternary cyclic codes, Signal Design and its Applications in Communications (IWSDA), 2015 seventh Intertional Workshop on Signal, DOI:10.1109/IWSDA.2015.7458415.

[11] C. Fan, N. Li, Z. Zhou, A class of optimal ternary cyclic codes and their duals, Finite Fields Appl. 37 (2016) 193-202.

[12] L. Wang, G. Wu, Several classes of optimal ternary cyclic codes with minimal distance four, Finite Fields Appl. 40 (2016) 126-137.

[13] P. Delsarte, On subfield subcodes of modified ReedCSolomon codes, IEEE Trans. Inf. Theory 21 (5) (1975) 575-576.

[14] C. Ding, Y. Yang, X. Tang, Optimal sets of frequency hopping sequences from linear cyclic codes, IEEE Trans. Inf. Theory 56 (7) (2010) 3605-3612.

[15] X. Liu, M. Harrison, Y. Luo, A note on the five valued conjectures of Johansen, Helleseth and Kholosha and zeta functions, IEEE Communication Letters 18 (9) (2014) 1483-1486.

[16] K. Feng, J. Luo, Value distribution of exponential sums from perfect nonlinear functions and their applications, IEEE Trans. Inf. Theory 53 (9) (2007) 3035-3041.

[17] T. Feng, On cyclic codes of length $2^{2^r} - 1$ with two zeros whose dual codes have three weights, Des. Codes Cryptogr. 62 (2012) 253-258.

[18] David M. Burton, Elementary Number Theory, Tata McGraw-Hill, 2006.

[19] W.C. Huffman, V. Pless, Fundamentals of Error-Correcting Codes, Cambridge University Press, Cambridge, 2003.

[20] T. Klove, Codes for Error Detection, World Scientific, 2007.

[21] C. Li, Q. Yue, F. Li, Weight distributions of cyclic codes with respect to pairwise coprime order elements, Finite Fields Appl. 28 (2014) 94-114.

[22] C. Li, N. Li, T. Helleseth, C. Ding, The weight distributions of several classes of cyclic codes from APN monomials, IEEE Trans. Inf. Theory 60 (8) (2014) 4710-4721.

[23] R. Lidl, H. Niederreiter, Finite Fields, Encycl. Math. Appl., vol.20 Addison-Wesley Reading, MA, 1983.

[24] B. Schmidt, C. White, All two-weight irreducible cyclic codes, Finite Fields Appl. 8 (2002) 1-17.

[25] J. Yang, M. Xiong, C. Ding, J. Luo, Weight distribution of a class of cyclic codes with arbitrary number of zeros, IEEE Trans. Inf. Theory 59 (9) (2013) 5985-5993.

[26] J. Yuan, C. Carlet, C. Ding, The weight distribution of a class of linear codes from perfect nonlinear functions, IEEE Trans. Inf. Theory 52 (2) (2006) 712-717.

[27] D. Zheng, X. Wang, H. Hu, X. Zeng, The weight distributions of two classes of p-ary cyclic codes, Finite Fields Appl. 29 (2014) 202-242.

[28] L. Liu, L. Li, X. Kai, S. Zhu, Repeated-root constacyclic codes of length $3lp^s$ and their dual codes, Finite Fields Appl. 42 (2016) 269-295.

[29] Z. Zhou, C. Ding, A class of three-weight cyclic codes, Finite Fields Appl. 25 (2014) 79-93.

[30] Z. Zhou, C. Ding, Seven classes of three-weight cyclic codes, IEEE Trans. Commun. 61 (10) (2013) 4120-4126.